



CRISPR
THERAPEUTICS

The logo consists of the word "CRISPR" in a bold, black, sans-serif font, positioned above the word "THERAPEUTICS" in a smaller, black, sans-serif font. Both words are contained within a black rectangular border. A small registered trademark symbol (®) is located at the bottom right of the border.



A photograph of a woman with a colorful patterned headscarf hugging a young girl from behind. The girl is smiling and holding a blue object. The background is blurred, showing an indoor setting. The image is overlaid with a semi-transparent white box containing text and a yellow bar at the bottom.

Creating transformative gene-based medicines for serious diseases

Corporate Overview | August 2021

Forward-Looking Statements



The presentation and other related materials may contain a number of “forward-looking statements” within the meaning of the Private Securities Litigation Reform Act of 1995, as amended, including statements regarding CRISPR Therapeutics’ expectations about any or all of the following: (i) the safety, efficacy and clinical progress of our various clinical programs; (ii) the status of clinical trials (including, without limitation, the expected timing of data releases, filing of clinical trial applications and INDs, any approvals thereof and the timing of commencement of clinical trials), development timelines and discussions with regulatory authorities related to product candidates under development by CRISPR Therapeutics and its collaborators; (iii) expectations regarding the data that is being presented; (iv) the number of patients that will be evaluated, the anticipated date by which enrollment will be completed and the data that will be generated by ongoing and planned clinical trials, and the ability to use that data for the design and initiation of further clinical trials; (v) the activities under CRISPR Therapeutics’ collaborations and the expected benefits thereof; (vi) the intellectual property coverage and positions of CRISPR Therapeutics, its licensors and third parties as well as the status and potential outcome of proceedings involving any such intellectual property; (vii) the sufficiency of CRISPR Therapeutics’ cash resources; and (viii) the therapeutic value, development, and commercial potential of CRISPR/Cas9 gene editing technologies and therapies. Without limiting the foregoing, the words “believes,” “anticipates,” “plans,” “expects” and similar expressions are intended to identify forward-looking statements. You are cautioned that forward-looking statements are inherently uncertain. Although CRISPR Therapeutics believes that such statements are based on reasonable assumptions within the bounds of its knowledge of its business and operations, forward-looking statements are neither promises nor guarantees and they are necessarily subject to a high degree of uncertainty and risk. Actual performance and results may differ materially from those projected or suggested in the forward-looking statements due to various risks and uncertainties. These risks and uncertainties include, among others: the potential for initial and preliminary data from any clinical trial and initial data from a limited number of patients not to be indicative of final trial results; the potential that the outcomes for each of CRISPR Therapeutics’ planned clinical trials and studies may not be favorable; that one or more of CRISPR Therapeutics’ internal or external product candidate programs will not proceed as planned for technical, scientific or commercial reasons; that future competitive or other market factors may adversely affect the commercial potential for CRISPR Therapeutics’ product candidates; uncertainties inherent in the initiation and completion of preclinical studies for CRISPR Therapeutics’ product candidates; availability and timing of results from preclinical studies; whether results from a preclinical trial will be predictive of future results of the future trials; uncertainties about regulatory approvals to conduct trials or to market products; CRISPR Therapeutics may not realize the potential benefits of its collaborations; potential impacts due to the coronavirus pandemic; uncertainties regarding the intellectual property protection for CRISPR Therapeutics’ technology and intellectual property belonging to third parties, and the outcome of proceedings (such as an interference, an opposition or a similar proceeding) involving all or any portion of such intellectual property; and those risks and uncertainties described under the heading “Risk Factors” in CRISPR Therapeutics’ most recent annual report on Form 10-K, quarterly report on Form 10-Q, and in any other subsequent filings made by CRISPR Therapeutics with the U.S. Securities and Exchange Commission, which are available on the SEC’s website at www.sec.gov. Existing and prospective investors are cautioned not to place undue reliance on these forward-looking statements, which speak only as of the date they are made. CRISPR Therapeutics disclaims any obligation or undertaking to update or revise any forward-looking statements contained in this presentation, other than to the extent required by law.

CRISPR THERAPEUTICS® standard character mark and design logo, CTX001™, CTX110™, CTX120™, and CTX130™ are trademarks and registered trademarks of CRISPR Therapeutics AG. All other trademarks and registered trademarks are the property of their respective owners.

CRISPR Therapeutics Highlights



Leading gene editing company focused on translating revolutionary CRISPR/Cas9 technology into transformative therapies



Achieving functional cures with CTX001™ in β -thalassemia and sickle cell disease



Next-generation immuno-oncology platform underlying wholly-owned, potentially best-in-class gene-edited allogeneic cell therapies CTX110™, CTX120™ and CTX130™



Enabling regenerative medicine 2.0 with CRISPR/Cas9-edited allogeneic stem cells



Advancing *in vivo* applications based on in-licensed technologies, platform improvement and strategic partnerships

The CRISPR/Cas9 Revolution

A **SPECIFIC**, **EFFICIENT** and **VERSATILE** tool for editing genes



Disrupt



Delete



Correct or Insert

*“If scientists can dream of a genetic manipulation,
CRISPR can now make it happen”*

Science

Our Pipeline




PROGRAM	RESEARCH	IND-ENABLING	CLINICAL	MARKETED	STATUS	PARTNER	STRUCTURE
 Hemoglobinopathies							
CTX001™: β-thalassemia					Enrolling		Collaboration
CTX001™: Sickle cell disease (SCD)					Enrolling		Collaboration
 Immuno-oncology							
CTX110™: Anti-CD19 allogeneic CAR-T					Enrolling		Wholly-owned
CTX120™: Anti-BCMA allogeneic CAR-T					Enrolling		Wholly-owned
CTX130™: Anti-CD70 allogeneic CAR-T					Enrolling		Wholly-owned
 Regenerative medicine							
Type I diabetes mellitus					PhI/II in 2021		Collaboration
 In vivo approaches							
Glycogen storage disease Ia (GSD Ia)							Wholly-owned
Duchenne muscular dystrophy (DMD)							License
Myotonic dystrophy type 1 (DM1)							Collaboration
Cystic fibrosis (CF)							License

Additional undisclosed, early-stage programs subject to collaboration or license agreements with Vertex and Bayer



Recent Collaborations to Combine Leading Capabilities




Gene editing
and T cells

+

Natural Killer
(NK) cells

Focus on co-developing and co-commercializing three gene-edited, donor-derived NK therapies:

- CAR-NK program targeting CD70
- Combined NK and T cell program (NK+T) to harness the synergies of the innate and adaptive immune systems
- CAR-NK program optioned from Nkarta's gene-edited pipeline



Gene
editing

+

Novel AAV
vectors

Focus on advancing gene-edited therapies for familial amyotrophic lateral sclerosis (ALS) and Friedreich's ataxia:

- CRISPR leads Friedreich's ataxia program, and performs gene-editing activities for both programs
- Capsida leads ALS program, and conducts capsid engineering and manufacturing for both programs
- CRISPR and Capsida each have the option to co-develop and co-commercialize the program that the other leads

Hemoglobinopathies – Devastating Blood Diseases

Sickle Cell Disease (SCD) and β -Thalassemia

Blood disorders caused by mutations
in the β -globin gene



Sickled



Normal Cell



Thalassemic

Significant worldwide burden

ANNUAL BIRTHS

300K

SCD



60K

β -thalassemia

High morbidity and mortality



Anemia




Pain



Early death


Heavy burden of patient care




Frequent transfusions and hospitalizations

Our Approach – Upregulating Fetal Hemoglobin

Symptoms in SCD and β -Thalassemia Decrease as HbF Level Increases



Powars, et al. Blood 1984



Musallam, et al. Blood 2012

- **Naturally occurring genetic variants** cause a condition known as **hereditary persistence of fetal hemoglobin (HPFH)**, which **leads to reduced or no symptoms** in patients with SCD and β -thalassemia
- **Our gene editing strategy aims to mimic these variants in symptomatic patients**, an approach supported by well-understood genetics

Pioneering CRISPR Trials



Design	Phase 1/2, international, multi-center, open-label, single arm studies to assess the safety and efficacy of CTX001 in patients with transfusion-dependent β -thalassemia (TDT) and SCD, respectively	
Target enrollment	45 patients aged 12 - 35 years with TDT, including β^0 / β^0 genotypes, defined as a history of at least 100 mL/kg/year or 10 units/year of pRBC transfusions in the previous 2 years	45 patients aged 12 - 35 years with severe SCD and a history of ≥ 2 vaso-occlusive crises/year over the previous two years
Primary endpoint	Proportion of patients achieving sustained transfusion reduction of 50% for at least 6 months starting 3 months after CTX001 infusion	Proportion of patients with HbF $\geq 20\%$, sustained for at least 3 months starting 6 months after CTX001 infusion

>45 patients with TDT and SCD dosed across both trials as of July 29, 2021

TDT: Patient Baseline and Treatment Characteristics

Patients with ≥ 3 -month follow-up (n=15)

Patient baseline

	n
Genotype	β^0 / β^0
	$\beta^0 / \text{IVS-I-110}^1$
	$\text{IVS-I-110} / \text{IVS-I-110}^1$
	β^0 / β^E
	β^0 / β^+
	β^+ / β^+
Gender	
Female/Male	9/6
	Median (range)
Age at consent	
Years	23 (18-32)
Pre-study pRBC transfusions ²	
Units/year	34 (20.5-61)

Treatment characteristics

	Median (range)
Drug product cell dose	
CD34+ cells x $10^6/\text{kg}$	6.5 (3.5-16.6)
Neutrophil engraftment ³	
Study day ⁴	29 (19-39)
Platelet engraftment ⁵	
Study day ⁴	40 (29-56)
Duration of follow-up	
Months	8.7 (4.0-26.2)

Data as of March 30, 2021

(1) IVS-I-110 phenotype is severe and similar to β^0 ; (2) Annualized number during the 2 years before consenting to study participation; (3) Defined as the first day of 3 measurements of absolute neutrophil count ≥ 500 cells/ μL on 3 consecutive days; (4) Study day 1 is the day of CTX001 infusion; (5) Defined as the first day of 3 consecutive measurements of platelet count $\geq 20,000/\mu\text{L}$ on 3 different days after CTX001 infusion, without a platelet transfusion in the past 7 days

TDT: Summary of Adverse Events

Patients with ≥ 3 -month follow-up (n=15)

Safety profile generally consistent with myeloablation and autologous stem cell transplant

	Patients with non-serious AEs, n	Patients with SAEs, n
Relationship ¹		
Related to plerixafor and/or G-CSF	10	0
Related to busulfan only	15	3
Related to CTX001 only	1 ²	1
Related to busulfan and CTX001	3 ³	1
Not related to any study drug	15	9

- 3 patients experienced SAEs assessed as related or possibly related to busulfan only: venoocclusive liver disease (2 patients), febrile neutropenia (1 patient), colitis (1 patient), and pneumonia (1 patient); all have resolved
- 1 patient had 4 SAEs assessed by the investigator as related or possibly related to CTX001: headache, haemophagocytic lymphohistiocytosis (HLH), acute respiratory distress syndrome, and idiopathic pneumonia syndrome (latter also related to busulfan); all occurred in the context of HLH and have resolved
- In addition to the above, one patient with <3 months of follow-up experienced an SAE of cerebellar hemorrhage, assessed by the investigator to be life-threatening, related to busulfan-induced thrombocytopenia, and not related to CTX001; the SAE has since resolved


Data as of March 30, 2021

(1) Includes related, possibly related, and missing relationship AEs; (2) 1 patient experienced a non-serious AE of anaemia possibly related to CTX001 (resolved); (3) 3 patients experienced non-serious AEs related or possibly related to busulfan and CTX001: petechiae, pyrexia, epistaxis, lymphocyte count decreased, neutrophil count decreased, WBC count decreased, and platelet count decreased (all resolved)

TDT: Clinically Meaningful HbF and Total Hb Are Achieved Early and Maintained


Hemoglobin fractionation, Hb (g/dL)

HbF HbA HbA2



Data as of March 30, 2021

TDT: Duration of Transfusion Independence After CTX001



Data as of March 30, 2021

(1) IVS-I-110 phenotype is severe and similar to β^0

SCD: Patient Baseline and Treatment Characteristics

Patients with ≥3-month follow-up (n=7)

Patient baseline

		n
Genotype	β^s / β^s	7
Gender	Female/Male	3/4
		Median (range)
Age at consent	Years	22 (19-34)
Pre-study VOCs	VOCs/year ¹	5.5 (2.5-9.5)

Treatment characteristics

	Median (range)
Drug product cell dose CD34+ cells x 10 ⁶ /kg	3.3 (3.1-3.9)
Neutrophil engraftment ² Study day ³	25 (17-33)
Platelet engraftment ⁴ Study day ³	33 (30-53)
Duration of follow-up Months	7.6 (4.9-22.4)

Data as of March 15, 2021

(1) Annualized rate during the 2 years before consenting to study participation; (2) Defined as the first day of 3 measurements of absolute neutrophil count ≥500 cells/μL on 3 consecutive days; (3) Study day 1 is the day of CTX001 infusion; (4) Defined as the first day of 3 consecutive measurements of platelet count ≥50,000/μL on 3 different days after CTX001 infusion, without a platelet transfusion in the past 7 days

SCD: Summary of Adverse Events

Patients with ≥ 3 -month follow-up (n=7)

Safety profile generally consistent with myeloablation and autologous stem cell transplant

	Patients with non-serious AEs, n	Patients with SAEs, n
Relationship ¹		
Related to plerixafor only	6	2
Related to busulfan only	7	1
Related to CTX001 only	0	0
Related to busulfan and CTX001	3 ²	0
Not related to any study drug	7	6


- Post-CTX001 infusion, 1 patient experienced an SAE related to busulfan of sepsis, which resolved
- No SAEs related to CTX001 were reported

Data as of March 15, 2021

(1) Includes related, possibly related, and missing relationship AEs; (2) 3 patients experienced non-serious AEs related or possibly related to busulfan and CTX001: dermatitis, lymphopenia, and CD4 lymphocytes decreased

SCD: Clinically Meaningful HbF and Total Hb Are Achieved Early and Maintained

Hemoglobin fractionation, Hb (g/dL)




Data as of March 15, 2021

SCD: Duration VOC-Free After CTX001

Pre-study VOC burden

Average number per year
over the previous 2 years



Improvements in markers of hemolysis (serum lactate dehydrogenase and haptoglobin) observed;
haptoglobin detectable by Month 6 in all 4 patients with Month 6 values


Pancellular HbF Expression and Durable Editing

TDT
SCD



Pancellular expression of HbF maintained

Mean % peripheral F-cells (*range*), % circulating RBCs expressing HbF



Data as of March 30, 2021 for TDT and March 15, 2021 for SCD

(1) Bone marrow editing assessments performed starting at 6 months, 12 months, and 24 months of follow-up

Allogeneic CAR-T Therapy Has Transformative Potential

Before Patient Diagnosis

Autologous: patient derived




PATIENT

Allogeneic: healthy-donor derived



HEALTHY DONOR



T Cells

Manufacture


100+ Doses

After Patient Diagnosis

WEEK 1

WEEK 2

WEEK 3



Apheresis



Manufacture



Single
Treatment




Treatment

- **Off-the-shelf:** Immediate treatment without risk of manufacturing failure, saving patients valuable time in which their disease could progress
- **Flexible dosing** (e.g., re-dosing)
- **A more consistent product**
- **Scalable manufacturing and simpler logistics**
- **Broader accessibility**

CTX110: Differentiated CRISPR-Edited Allogeneic CAR-T Design

Multiplex CRISPR gene editing in one step designed to:

- **Improve persistence in the allo setting** via $\beta 2M$ knock-out to eliminate MHC I expression
- **Avoid need** for more toxic lymphodepletion regimens




- **Prevent GvHD** via TCR disruption
- **Improve consistency and safety by precise insertion** of CAR construct into *TRAC* locus without using lentivirus or retrovirus

*CTX120TM and CTX130TM utilize the **same CRISPR-edited allogeneic T cell design**, but with different CAR targets, as well as additional editing in the case of CTX130*

CARBON: Trial Design

CARBON: Single-arm study evaluating the safety and efficacy of CTX110

Allogeneic CAR-T enables simplified trial design: short screening timeframe, no apheresis, no bridging chemotherapy, and on-site availability of CAR-T cell product



Key eligibility criteria

- Age ≥18 years
- Relapsed/refractory non-Hodgkin lymphoma, as evidenced by 2+ lines of prior therapy
- ECOG performance status 0 or 1
- Adequate renal, liver, cardiac, and pulmonary organ function
- No prior allogeneic SCT or treatment with CAR-T therapy

Primary endpoints


- Incidence of adverse events, defined as DLTs
- ORR

Key secondary endpoints

- DoR, PFS, and OS

CARBON: Patient Flow and Baseline Characteristics

As of the data cutoff date:



N (%) (unless otherwise noted)

Cell dose (CAR ⁺ T cells)	DL1 30x10 ⁶ N=3	DL2 100x10 ⁶ N=3	DL3 300x10 ⁶ N=4	DL4 600x10 ⁶ N=1
Median age, years (range)	52 (50-61)	64 (58-74)	64.5 (62-74)	72
Male	2 (66.7)	2 (66.7)	1 (25)	1 (100)
Lymphoma subtypes				
Diffuse large B-cell lymphoma (DLBCL) ¹	3 (100)	3 (100)	4 (100)	1 (100)
Follicular lymphoma	0	0	0	0
Current disease stage (per Lugano 2014)²				
Stage III	1 (33.3)	1 (33.3)	2 (50)	0
Stage IV	2 (66.7)	2 (66.7)	1 (25)	1 (100)
Prior treatments				
Median number (range)	2.0 (2-8)	3.0 (2-3)	2.0 (2-4)	5
Hematopoietic stem cell transplant	0	0	3 (75)	1 (100)
Refractory to last therapy	3 (100)	3 (100)	0	0

(1) Including high grade lymphoma (e.g., triple hit), transformed follicular lymphoma (tFL), Richter's Transformation; (2) One patient with Stage II disease treated at DL3

Dose-Dependent Responses Observed with CTX110


Best response per 2014 Lugano criteria¹ by independent central assessment

Cell dose (CAR+ T cells)	DL1 30x10 ⁶ N=3	DL2 100x10 ⁶ N=3	DL3 300x10 ⁶ N=4	DL4 600x10 ⁶ N=1
Overall response rate (ORR), N (%)	0 (0%)	1 (33%)	2 (50%)	1 (100%)
Complete response (CR) rate, N (%)	0 (0%)	1 (33%)	2 (50%)	1 (100%)

- Early evidence of **dose response, with complete responses achieved in 4 patients**
- **Responses achieved without the use of more toxic lymphodepletion agents**, consistent with engineering of CTX110 for immune evasion
- **CAR-T cells detected at multiple time points in all patients** in DL2-4, with **consistent peak expansion** of CTX110 in the peripheral blood seen around 1-2 weeks post infusion and CTX110 detected out **as late as 180 days** after administration

Dose-Dependent Reduction in Tumor Size with CTX110


Best tumor size reduction per 2014 Lugano criteria by independent central assessment



* Patient subsequently failed autologous CAR-T

Data as of September 28, 2020

Complete Responses with CTX110 Showed Durability at Month 3 and Beyond



Imaging per protocol occurs at M1, M3, and M6; * Patient died while in CR at Day 52 post CTX110 infusion following data cutoff

Data as of September 28, 2020

Acceptable Safety Profile with CTX110 at DL3 and Below

Treatment-emergent adverse events (AEs) of special interest in DL1-3, N (%)

N=10	Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
Graft-versus-Host Disease (GvHD)	0	0	0	0	0
Cytokine Release Syndrome (CRS) ^{1,2}	1 (10%)	2 (20%)	0	0	0
ICANS ^{1,3}	0	1 (10%)	0	0	0
Infections	0	0	1 (10%)	0	0

For patients in DL1 through DL3 (N=10):

- **No GvHD** despite all patients with $\leq 3/12$ HLA match to CTX110 donors
- **No CRS or ICANS above Grade 2**
- **No infusion reactions**
- **4 serious adverse events (SAEs) following CTX110 infusion not related to disease progression among 3 treated patients:** ICANS (n=1), CRS (n=1), periorbital cellulitis (n=1), febrile neutropenia (n=1)

Safety for patient treated at DL4 (600x10⁶ CAR⁺ T cells):

- Patient had received five prior lines of therapy, including autologous stem cell transplant
- Experienced Grade 2 CRS at Day 5 that resolved
- Admitted with febrile neutropenia at Day 26 and developed confusion and memory loss starting at Day 28, with further deterioration ultimately requiring intubation for airway protection
- Initially treated for ICANS and later found to have reactivation of HHV-6 and HHV-6 encephalitis
- Despite treatments, patient remained obtunded and died on Day 52 after family requested withdrawal of care

(1) Per ASTCT criteria; other AEs graded per CTCAE; (2) Includes two separate episodes of CRS (1 G1, 1 G2) in single patient; worst grade reported;

(3) Immune effector Cell-Associated Neurotoxicity Syndrome

Our I/O Strategy and Allogeneic CAR-T Pipeline



	PROGRAM	RESEARCH	IND-ENABLING	CLINICAL	MARKETED	STATUS
Validate <i>allogeneic platform with proven targets</i>	CTX110 (anti-CD19) <i>B-cell malignancies</i>					<i>Enrolling</i>
	CTX120 (anti-BCMA) <i>Multiple myeloma</i>					<i>Enrolling</i>
Expand <i>from hematologic cancers into solid tumors</i>	CTX130 (anti-CD70) <i>T- and B-cell lymphomas</i>					<i>Enrolling</i>
	CTX130 (anti-CD70) <i>Renal cell carcinoma</i>					<i>Enrolling</i>
Unlock <i>the full potential of I/O cell therapy with CRISPR</i>	Anti-CD33 allogeneic CAR-T					<i>Incorporating additional editing, novel targeting, etc.</i>
	Anti-PTK7 allogeneic CAR-T					
	Additional undisclosed programs					


CRISPR Enables the Next Generation of I/O Cell Therapy



CRISPR gene editing facilitates consistent, multiplex editing to:

- Produce allogeneic cell therapies
- Enhance immune cell performance
- Speed the discovery and generation of novel therapeutic candidates


Multiplexed, single-shot 6x knock-out plus CAR insertion performed at high efficiency



6x-edited CAR-T cells show no viability decrease, no cytokine-independent growth and robust target-specific cytotoxicity

CRISPR Enables Regenerative Medicine 2.0

CRISPR/Cas9 Technology Opens Broader Applications for Regenerative Medicine



Unlocking *In Vivo* Applications of CRISPR/Cas9

AAV Vectors for Neuromuscular Indications

- **Adeno-associated virus (AAV)** to deliver Cas9 and gRNA to muscle, the nervous system and other tissues
- Collaboration with StrideBio to improve tissue specificity and reduce immunogenicity
- Programs include DMD and DM1 in collaboration with Vertex, as well as other early research programs



LNPs for Liver Indications

- **Lipid nanoparticles (LNPs)** containing mRNA encoding Cas9 and gRNA for delivery to the liver
- mRNA technology from CureVac
- Programs include GSD Ia and other early research programs

Enabling collaborations



Optimizing the CRISPR/Cas9 Platform


Nuclease Engineering

Enhance CRISPR/Cas9 system through protein engineering



Guide RNA Optimization

Identify optimal guide RNA formats and sequences for therapeutic editing




Advanced Editing

Improve efficiency of gene correction and multiplexing



Synthetic Biology

Engineer improved cellular therapeutics



Strong U.S. and Global Foundational IP Position



United States

Charpentier / UC Berkeley / U. Vienna granted patents of broad scope; multiple applications progressing

40

Patents of broad scope granted, including the patent involved in the 1st interference

36+

Additional patent applications moving forward in parallel with both broad and narrow claims, including 4 patent applications of broad scope allowed

2nd

Interference with Broad Institute in priority phase to determine who was first to invent CRISPR/Cas9 gene editing in eukaryotic cells; separate interference declared with Toolgen on same subject matter



Europe and Global

Charpentier / UC Berkeley / U. Vienna granted foundational patents, including use in eukaryotes

3

Patents of broad scope granted in the EU

31

Patents of broad scope granted in the UK, Germany, Japan, China, Singapore, Hong Kong, Ukraine, Israel, Australia, New Zealand, Mexico, South Africa and elsewhere

~80

Jurisdictions worldwide in which applications with both broad and narrow claims are advancing

Building a Great Company



EXPERIENCED
Management Team

**END-TO-END
CAPABILITIES**
With >400 Employees

**COLLABORATIVE &
ENTREPRENEURIAL**
Culture